



EUROPEAN
HEMATOLOGY
ASSOCIATION

EHA25 VIRTUAL

25th Congress of the European Hematology Association

VIRTUAL EDITION

Initial Results of a Phase 2 Study of Loncastuximab Tesirine, a Novel Pyrrolobenzodiazepine-Based Antibody-Drug Conjugate, in Patients With Relapsed or Refractory Diffuse Large B-Cell Lymphoma

Carmelo Carlo-Stella, MD¹, Pier Luigi Zinzani, MD², Brad Kahl, MD³, Paolo F. Caimi, MD⁴, Melhem Solh, MD⁵, Kirit Ardeshta, MD⁶, Anastasios Stathis, MD⁷, Mehdi Hamadani, MD⁸, Weiyun Ai, MD, PhD⁹, Brian Hess, MD¹⁰, Juan Pablo Alderuccio, MD¹¹, Jay Feingold, MD, PhD¹², David Ungar, MD¹², Shui He, PhD¹², Yajuan Qin, MD, PhD¹², John Radford, MD, FMedSci¹³

¹Department of Oncology and Hematology, Humanitas Cancer Center, Humanitas University, Milan, Italy; ²Institute of Hematology "Seràgnoli" University of Bologna, Bologna, Italy; ³Department of Medicine, Oncology Division, Washington University, St. Louis, MO, USA; ⁴University Hospitals Cleveland Medical Center/Case Western Reserve University, Cleveland, OH, USA; ⁵Blood and Marrow Transplant Program at Northside Hospital, Atlanta, GA, USA; ⁶Department of Hematology, University College London Hospitals NHS Foundation Trust, London, UK; ⁷Oncology Institute of Southern Switzerland, Bellinzona, Switzerland; ⁸Division of Hematology and Oncology, Medical College of Wisconsin, Milwaukee, WI, USA; ⁹Division of Hematology and Oncology, Department of Medicine, University of California, San Francisco, CA, USA; ¹⁰Division of Hematology and Medical Oncology, Department of Medicine, Medical University of South Carolina, Charleston, SC, USA; ¹¹Sylvester Comprehensive Cancer Center, University of Miami, Miami, FL, USA; ¹²ADC Therapeutics America Inc., Murray Hill, NJ, USA; ¹³University of Manchester and The Christie NHS Foundation Trust, Manchester Academic Health Centre, Manchester, UK

Disclosures and Acknowledgements

C. Carlo-Stella: research support from ADC Therapeutics, Sanofi, Roche; consultant or advisor for Boehringer Ingelheim, Sanofi, ADC Therapeutics, Servier, Roche, Genenta Science; honoraria from Bristol-Myers Squibb, Merck Sharp & Dohme, Takeda, Novartis, Janssen.

P. L. Zinzani: consultant to Verastem, MSD, Sanofi and Eusapharma; speaker for Verastem, Celltrion, Gilead, Janssen-Cilag, BMS, Servier, MSD, Immune Design, Celgene, Portola, Roche, Eusapharma, Kyowa Kirin; advisory boards for Verastem, Celltrion, Gilead, Janssen-Cilag, BMS, Servier, Sandoz, MSD, Immune Design, Celgene, Portola, Roche, Eusapharma, Kyowa Kirin, ADC Therapeutics; research support from ADC Therapeutics.

B. Kahl: research support from ADC Therapeutics; advisor or consultant for ADC Therapeutics, Seattle Genetics, Genentech.

P. F. Caimi: research support from ADC Therapeutics; advisory boards for ADC Therapeutics, Genentech, Kite Pharma; speaker for Celgene.

M. Solh: research support from ADC Therapeutics; speaker bureau for Amgen, Celgene.

K. Ardeschna: supported by the University College London (UCL)/UCL Hospitals (UCLH) Biomedical Research Unit; honoraria/advisory boards for ADC Therapeutics, Celgene, Gilead, Takeda, Roche, Beigene.

A. Stathis: institutional research funding from Roche, Pfizer, Merck, Bayer, Novartis, MEI-Pharma, ADC Therapeutics; other remuneration from Abbvie, PharmaMar.

M. Hamadani: research support from ADC Therapeutics, Otsuka, Takeda, Sanofi Genzyme, MedImmune, Merck; speaker bureau for Sanofi Genzyme; advisor/consultant for MedImmune, Jansen, Celgene, ADC Therapeutics.

W. Ai: research support from Nurix Therapeutics; advisory boards for ADC Therapeutics, Nurix, Kymera.

B. Hess: no relevant disclosures.

J. P. Alderuccio: honoraria from OncLive and Oncinfo; immediate family member has served on advisory boards for Puma Biotechnology, Inovio Pharmaceuticals, Agios Pharmaceuticals, Forma Therapeutics and Foundation Medicine.

J. Feingold, D. Ungar, S. He, and Y. Qin: employees of ADC Therapeutics with stock options.

J. Radford: consultant/advisory role for Takeda, ADC Therapeutics, Bristol-Myers Squibb, Novartis, Kite Pharma; speaker for Takeda, ADC Therapeutics, Seattle Genetics; research support from Takeda; provided expert testimony for Takeda and ADC Therapeutics; stocks/shares: AstraZeneca and GSK (spouse).

This study was funded by ADC Therapeutics SA (NCT03589469)

Acknowledgements

The authors would like to thank all the participating patients and their families, all study co-investigators and research coordinators.


Editorial support was provided by Louise Gildea, PhD, and Becky Salisbury, PhD, at Fishawack Health, funded by ADC Therapeutics.

Loncastuximab Tesirine (ADCT-402)

Despite recent advances in DLBCL treatment, outcomes for patients with R/R DLBCL remain poor¹

- Lonca is an ADC comprising a humanised anti-CD19 antibody conjugated to a potent PBD dimer²
- Lonca had encouraging antitumour activity and acceptable safety in R/R DLBCL in a Phase 1 first-in-human trial³

Lonca targets CD19, which is expressed in the majority of B-cell malignancies⁴



1. Lonca binds to CD19 antigen and is internalised
2. The linker is cleaved and PBD dimers released
3. Cytotoxic DNA cross-links are formed
4. The DNA replication fork stalls
5. The cell goes into apoptosis


Study Design: Single-arm, Open-label Phase 2 Study

Patient population:

Patients with R/R DLBCL following ≥ 2 lines of prior systemic therapy

Primary objective:

Evaluate efficacy, using ORR (central review), and safety of the full Phase 2 study population



Futility requirements met:
ORR for first 52 patients¹



Total enrolment:
145 patients

1. Carlo-Stella C, *et al. Blood*. 2019;134(Supp1):757. DLBCL, diffuse large B-cell lymphoma; Lonca, loncastuximab tesirine; ORR, overall response rate; Q3W, every 3 weeks; Q12W, every 12 weeks; R/R, relapsed/refractory.


Baseline Characteristics

Patient characteristics		Total (N=145)
Sex, n (%)	Female	60 (41.4)
	Male	85 (58.6)
Age, years, median (min, max)		66.0 (23–94)
Histology, n (%)	DLBCL	127 (87.6)
	HGBCL	11 (7.6)
	PMBCL	7 (4.8)
Double/triple hit, n (%)		15 (10.3)
Double/triple expressor, n (%)		20 (13.8)
Transformed disease, n (%)		29 (20.0)
Stage, n (%)	I–II	33 (22.8)
	III–IV	112 (77.2)

Patient treatment history		Total (N=145)
No. of previous systemic therapies,* median (range)		3 (2–7)
First-line systemic therapy response, n (%)	Relapse	99 (68.3)
	Refractory†	29 (20.0)
	Other‡	17 (11.7)
Last-line systemic therapy response,¶ n (%)	Relapse	43 (29.7)
	Refractory†	84 (57.9)
	Other‡	18 (12.4)
Refractory to all prior therapies, n (%)	Yes	25 (17.2)
	No	115 (79.3)
	Other‡	5 (3.4)
Prior stem cell transplant, n (%)	Allogeneic	2 (1.4)
	Autologous	21 (14.5)
	Both	1 (0.7)

145 patients were enrolled and received a mean of 4.3 cycles of Lonca (range: 1–15)



Response to Lonca by Histology




ORR in the total population was 48.3% (95% CI: 39.9, 56.7) and an additional 15.2% (22 pts) had stable disease

Subgroup Analysis of ORR

Patients refractory to first-line or last-line prior therapy had ORRs (CRR) of 37.9% (17.2%) and 36.9% (11.9%), respectively



Duration of Response



Median duration of response was 10.25 months (95% CI: 5.98, –)

TEAEs of Any Grade in $\geq 20\%$ of Patients (Safety Analysis Set; N=145)

Preferred term n (%)	Patients n (%)
Patients with any TEAE	143 (98.6%)
GGT increased	59 (40.7)
Neutropenia	57 (39.3)
Thrombocytopenia	48 (33.1)
Fatigue	40 (27.6)
Anaemia	38 (26.2)
Nausea	34 (23.4)
Cough	32 (22.1)
Alkaline phosphatase increased	29 (20.0)
Peripheral oedema	29 (20.0)

The most common grade ≥ 3 TEAEs ($\geq 10\%$ of patients) were:

- Neutropenia (37 patients; 25.5%)
 - Incidence of febrile neutropenia was low (5 patients; 3.4%)
- Thrombocytopenia (26 patients; 17.9%)
- GGT increased (24 patients; 16.6%)
- Anaemia (15 patients; 10.3%)

Conclusions

Lonca had substantial single-agent antitumour activity in patients with R/R DLBCL who failed established therapies

- ORR was 48.3% (70/145 patients; 95% CI: 39.9, 56.7)
- CRR was 24.1% (35/145 patients; 95% CI: 17.4, 31.9)
- Median duration of response was 10.25 months (95% CI: 5.98, –)
- The toxicity profile was manageable and no new safety concerns were identified
- There is the potential for Lonca to fill a critical unmet need for treatment of heavily pre-treated patients with DLBCL